

**WHAT IS CLAIMED IS:**

1. A method of determining an individual's predisposition to epilepsy and/or development of epilepsy, as well as  
5 predicting this individual's response to medication, said method comprising the step of determining the genotype of at least one gene selected from SCN1A, SCN2A and SCN3A of the individual, or of a DNA variant, equivalent, or mutation which shows a linkage disequilibrium therewith, thereby determining an individual's predisposition to epilepsy  
10 and/or development of epilepsy.
2. The method of claim 1, wherein the step of determining the SCN1A, SCN2A or SCN3A genotype comprises restriction endonuclease digestion.
3. The method of claim 1, wherein the step of  
15 determining the SCN1A, SCN2A or SCN3A genotype comprises hybridizing with allele specific oligonucleotides.
4. The method of claim 1, which further comprises a step, prior to determining the SCN1A, SCN2A or SCN3A genotype, of amplifying a segment of the the SCN1A, SCN2A or SCN3A using  
20 polymerase chain reaction.
5. The method of claim 1, wherein the step of determining the SCN1A, SCN2A or SCN3A genotype comprises a sequencing of SCN1A, SCN2A or SCN3A , or parts thereof.
6. The method of claim 1, wherein the SCN1A,  
25 SCN2A or SCN3A genotype is determined using a polymorphic variant

site in linkage disequilibrium with at least one allelic variant or mutant identified in accordance with the present invention.

7. An assay for screening a test agent and selecting an agent which modulates inactivation of a sodium channel involved in epilepsy comprising:

- a) a recombinant SCN1A, SCN2A or SCN3A gene which encodes an alpha subunit of said sodium channel or functional fragment thereof; and
  - b) assaying a function of said sodium channel;
- wherein an agent can be selected when an observable difference is observed between the inactivation of said sodium channel in the presence of said test agent, as compared to in an absence thereof, and wherein a malfunction of said sodium channel is associated with epilepsy.

8. An assay for screening a test agent and selecting an agent which modulates the activity of a sodium channel involved in epilepsy comprising:

- a) a recombinant SCN1A, SCN2A or SCN3A gene which encodes an alpha subunit of said sodium channel or functional fragment thereof; and
  - b) assaying the activity of said sodium channel;
- wherein an agent can be selected when an observable difference is observed between the activity of said sodium channel in the presence of said test agent, as compared to in an absence thereof, and wherein a malfunction of said sodium channel is associated with epilepsy.

9. A method of using specific alleles of the SCN1A, SCN2A or SCN3A genes, or a variant, equivalent, or mutation thereof which shows linkage disequilibrium therewith, to set-up a screening assay

for agents destined to modulate sodium channel function for the purpose of identifying agents having an application in epilepsy therapy.

5 10. A method for identifying, from a library of compounds, a compound with therapeutic effect on epilepsy or other neurological disorders comprising:

- 10 a) providing a screening assay comprising a measurable biological activity of SCN1A, SCN2A or SCN3A protein or gene;
- 10 b) contacting said screening assay with a test compound; and
- 10 c) detecting if said test compound modulates the biological activity of SCN1A, SCN2A or SCN3A protein or gene;

wherein a test compound which modulates said biological activity is a compound with said therapeutic effect.

15

11. The method of claim 10, wherein the test compound with said therapeutic effect is further modified by combinatorial or medicinal chemistry to provide further analogs of said test compound also having said therapeutic effect.

20

12. A compound having therapeutic effect on epilepsy or other neurological disorders, identified by a method comprising,

- 25 a) providing a screening assay comprising a measurable biological activity of SCN1A, SCN2A or SCN3A protein or gene;
- b) contacting said screening assay with a test compound; and
- c) detecting if said test compound modulates the biological activity of SCN1A, SCN2A or SCN3A protein or gene;

wherein a test compound which modulates said biological activity is a compound with said therapeutic effect.

- 5           13. The compound of claim 12, wherein the compound with said therapeutic effect is further modified by combinatorial or medicinal chemistry to provide analogs of said compound also having said therapeutic effect.

10  
15  
20  
25  
30  
35  
40  
45  
50  
55  
60  
65  
70  
75  
80  
85  
90  
95  
100  
105  
110  
115  
120  
125  
130  
135  
140  
145  
150  
155  
160  
165  
170  
175  
180  
185  
190  
195  
200  
205  
210  
215  
220  
225  
230  
235  
240  
245  
250  
255  
260  
265  
270  
275  
280  
285  
290  
295  
300  
305  
310  
315  
320  
325  
330  
335  
340  
345  
350  
355  
360  
365  
370  
375  
380  
385  
390  
395  
400  
405  
410  
415  
420  
425  
430  
435  
440  
445  
450  
455  
460  
465  
470  
475  
480  
485  
490  
495  
500  
505  
510  
515  
520  
525  
530  
535  
540  
545  
550  
555  
560  
565  
570  
575  
580  
585  
590  
595  
600  
605  
610  
615  
620  
625  
630  
635  
640  
645  
650  
655  
660  
665  
670  
675  
680  
685  
690  
695  
700  
705  
710  
715  
720  
725  
730  
735  
740  
745  
750  
755  
760  
765  
770  
775  
780  
785  
790  
795  
800  
805  
810  
815  
820  
825  
830  
835  
840  
845  
850  
855  
860  
865  
870  
875  
880  
885  
890  
895  
900  
905  
910  
915  
920  
925  
930  
935  
940  
945  
950  
955  
960  
965  
970  
975  
980  
985  
990  
995